



Frank J. Totta, D.O., A.G.A.F. Mary Lynne Lucido, M.D. Todd W. Kilgore, M.D. R. Wade McCullough, D.O. Benjamin R. Alsop, M.D. Chase M. Zeilenga, D.O. Amy Waller, A.P.R.N., B.C. Andrea Gray, A.P.R.N., B.C. Emily McClanahan, FNP-C Maddison Selley, FNP-C

110 NE Saint Luke's Blvd • Ste 530 • Lee's Summit, MO 64086 • 816-554-3838

fax: 816-554-1634 www.summitgi.com

EGD PREPARATION

| Patient | LOCATIONS: | |
|----------------|---|---|
| Doctor | St. Luke's Outpatient Surgery Center 120 NE St. Luke's Blvd. Lee's Summit, MO 64086 | Lee's Summit Medical Center 2100 SE Blue Pkwy. Lee's Summit, MO 64063 |
| Date | St. Luke's East Hospital | College Park (CP Surgicenter) |
| Arrival Time | 100 NE St. Luke's Blvd. Lee's Summit, MO 64086 | 10787 Nall Ave Ste 100 Overland Park, KS 66211 |
| Procedure Time | | |

To ensure that your test is accurate and complete, you <u>must</u> follow these directions. If you have questions, please call our office at 816-554-3838. Summit Gastroenterology is dedicated to providing you with quality health care. *You matter to us!*

INSTRUCTIONS

A week before your procedure:

Check with your prescribing doctor to make sure it is safe to hold prescription blood thinner:

days prior. No aspirin or plavix the day of the procedure.

- Nothing to eat or drink after midnight for procedures scheduled before noon.
- However, if your test is scheduled after 12 p.m., you may have water 6 hours prior to scope time.
- Hold prescription diet pills, ie-phentermine, 7 days prior to procedure.
- Hold weekly dosing of GLP's: Trulicity (Dulaglutide), Bydureon BCise (Exenatide), Mounjaro, Zepbound (Tirzepatide), Ozempic, Wegovy (Semaglutide). *HOLD 1 WEEK PRIOR TO PROCEDURE.
- Hold Steglatro (Ertugliflozin), 4 days prior to procedure. *HOLD FOR 96 HOURS PRIOR TO PROCEDURE.
- Hold Jardiance (empagliflozin), Invokanna (canagliflozin), Farxiga (dapagliflozin), 3 days prior to procedure. *HOLD FOR 72 HOURS PRIOR TO PROCEDURE.
- You will need a driver sedation is given someone will need to stay with you for 12 hours after procedure
- No work, no driving, operating machinery, or making legal decisions x 24 hours after procedure
- Take any heart, blood pressure, or seizure medications in the morning the day of the procedure with sip of water.
- Hold all other medications the day of the procedure.
- Hold insulin in the morning the day of the procedure until after procedure.
- No smoking or vaping the day of the procedure. No chewing tobacco. Avoid ALL tobacco products day of procedure. No chewing gum the day of the procedure.
- Hold daily dosing GLP's: Byetta (Exenatide), Saxenda (Liraglutide), Victoza (Liraglutide), Adlyxin (Lixisenatide), Rybelsus (Semaglutide). *HOLD DAY BEFORE AND DAY OF PROCEDURE.

* During your procedure you will be administered either general anesthesia or heavy sedation. For your safety, it is essential you follow the above listed instructions. * *

Failure to comply with these instructions will result in the cancellation or delay of your procedure. Judgment will be made by your anesthesia provider.

Cancellation Policy

\$100 fee for cancellation of endoscopy appointments with less than 72 hours notification.

Insurance Information

Our office may verify insurance benefits and contact you to discuss our procedure fee.